

Kentish side, extending from Blackheath to Erith, and attaining an elevation at Shooter's Hill of over 400 feet; subsoil, sand, gravel, overlying chalk.

Anchusa arvensis.
Artemisia vulgaris.
Bryonia dioica.
Calamintha Clinopodium.
Carex depauperata (?) *olim.*
Centaurea Scabiosa; *nigra.*
Circea lutetiana.
Clematis Vitalba.
Cornus sanguinea.
Daucus Carota.
Diplotaxis tenuifolia.
Echium vulgare.
Erigeron acris.
Erythraea Centaurium.
Euphorbia amygdaloides.
Filago germanica.
Galium verum; *Mollugo.*
Geranium dissectum.
Geum urbanum.
Heracleum Sphondylium.
Hypericum perforatum; *hirsutum.*
Inula Conyza.
Lactuca Scariola (?) (*sandpits*).
Lathyrus pratensis.

Lepidium campestre.
Ligustrum vulgare.
Linaria vulgaris.
Lotus corniculatus.
Medicago maculata; *lupulina.*
Mercurialis perennis.
Ononis arvensis.
Origanum vulgare.
Pastinaca sativa.
Prenanthes muralis.
Reseda lutea; *Luteola.*
Sanicula europæa.
Silene inflata.
Solidago Virgaurea.
Spergula arvensis.
Spergularia rubra.
Stachys sylvatica.
Tanacetum vulgare.
Torilis Anthriscus.
Trifolium arvense.
Tussilago Farfara.
Verbena officinalis.
Vicia Cracca; *sepium*; *tetrasperma.*
Etc., etc.

SHOOTER'S HILL COMMON.

Castle Wood is enclosed; but the common fronting Blackheath is still open to the public. There are no trees: scrub or brushwood only, of Birch, Hazel, Oak, Sallows, White-thorn, Black-thorn, Briars, Furze, Broom, and Bramble. Subsoil, gravel.

**Achillea Ptarmica.*
 **Aira flexuosa*; **cæspitosa.*
 **Betonica officinalis.*
 **Carex hirta*; **ovalis.*
 **Centaurea nigra.*
 **Hieracium vulgatum*; **umbellatum.*

**Lotus major*; **corniculatus.*
 **Ornithopus perpusillus.*
 **Potentilla Tormentilla*; **Fragariastrum*
 **repens, etc.*
 **Senecio sylvaticus.*
 **Solidago Virgaurea.*
 **Teucrium Scorodonia.*

11. WANDSWORTH COMMON.

An extensive open space, of nearly three miles in circumference, partly bordered with villas, &c., and traversed by the Crystal Palace Railway, which has exercised upon it the ordinary effects of drainage. It is more or less covered with Furze, Bracken, and Ling; some parts are in fact densely furze-grown; others entirely free from it. There are several small ponds and pits, whence gravel has been excavated; and the subsoil is of this material. Nothing out of the common is to be found here now, but formerly its speciality was *Stratiotes aloides*.

**Acetas Acetosella.*
 **Achillea Ptarmica*; *Millefolium.*

**Aira præcox.*
 **Alisma Plantago.*

**Alopecurus fulvus*.
 **Anthemis nobilis*.
 **Anthoxanthum odoratum*.
 **Betonica officinalis*.
 **Bidens tripartita*.
 **Campanula rotundifolia*.
 Camelina sativa (near New Wandsworth station).
 **Cardamine pratensis*.
 **Carduus acaulis*; **pratensis*.
 **Carex flava*; **hirta*; **glauca*; **ovalis*.
 **Centaurea nigra*.
 **Cerastium semidecandrum*.
 **Convolvulus arvensis*.
 **Elodea canadensis*.
 **Erigeron canadensis*.
 **Euphrasia officinalis*.
 **Filago minima*.
 **Galium saxatile*; **uliginosum*; **palustre*.
 **Geranium molle*; **Robertianum*.
 **Glyceria fluitans*.
 **Gnaphalium uliginosum*.
 **Helosciadium inundatum*.
 **Hieracium Pilosella*.
 **Hydrocotyle vulgaris*.
 **Hypericum humifusum*.
 **Hypochoeris radicata*.
 **Juncus effusus*; **glomeratus*; **supinus*.
 **Leontodon hirtus*; **hispidus*; **autumnalis*.
 **Lotus corniculatus*.
 **Lychnis diurna*.
 **Medicago maculata*.
 **Mentha Pulegioides*; **hirsuta*.

**Moenchia erecta*.
 **Montia fontana*.
 **Myosotis arvensis*; **palustris*.
 **Nardus stricta*.
 **Nasturtium officinale*; **terrestre*.
 **Ononis spinosa*.
 **Ornithopus perpusillus*.
 **Pedicularis sylvatica*.
 **Peplis Portula*.
 **Polygonum Hydropiper*, **Persicaria*.
 **Potamogeton natans*; **crispum*; **pussillum*.
 **Potentilla Tormentilla*; **reptans*; **anserina*.
 **Prunella vulgaris*.
 **Ranunculus aquatilis* (agg.); **Flammula*; **hederacea*; **sceleratus*; **acris*; **bulbosus*.
 **Sagina procumbens*.
 **Salix repens*.
 **Scabiosa succisa*.
 **Scirpus palustris*; **setaceus*; **fluitans*.
 **Senebiera Coronopus*.
 **Senecio Jacobaea*.
 **Trifolium repens*; **filiforme*.
 **Veronica hederifolia*; **arvensis*; **serpyllifolia*; **Chamaedrys*; **Beccabunga*.

CRYPTOGAMS.

**Bryum erythrocarpum*.
 **Ceratodon purpureus*.
 **Equisetum limosum*; **palustre*; **arvense*.
 **Pteris aquilina*.

Clapham Common is furzy in some parts, and has some fine groups of trees upon it; it is surrounded by houses. *Limosella aquatica* grew formerly on the margins of the central pond; may be there still. Tooting Common is two miles in circumference, and furze-grown in parts, with Briars and a few Oak-trees at the further extremity. **Dicranum scoparium*; **Polytrichum juniperinum*; **Cratægus Oxyacantha*, &c. Streatham Common is of the same character, but less extensive. **Anthemis nobilis*; **Trifolium subterraneum*; **Arenaria trinervis*; **Achillea Ptarmica*, &c.; on tombstones in Streatham churchyard, **Hypnum confertum*; **serpens*. They have both a gravelly subsoil. About Clapham Junction, **Erysimum cheiranthoides* (abundant). Near the station, Wandsworth, *Isatis tinctoria* has been lately observed. Streatham Common is separated from that of Tooting by an avenue of Elms. Its subsoil is the same, and elevated at the upper end, where there is some Furze and Bramble, over three hundred feet. *Senecio viscosus* grew here formerly; and in the closes, about, *Bunium flexuosum*; *Lactuca virosa*; *Daphne Laureola*, &c.

12. ROADSIDES, COPSES, AND WASTE PLACES ABOUT NORWOOD.

The low hills which constitute the Southern Heights of London attain their highest elevation at Norwood, in connection with which are Gipsy Hill, Sydenham Hill and Forest Hill. Norwood is under four hundred feet